

## Sampling methods

### Sampling from an unnormalized distribution

Let  $X$  be a random variable, of which the density  $f_X$  is known up to a normalizing constant :  $f_X(x) = Z_X^{-1}f(x)$  with  $Z_X$  unknown and  $f(x)$  defined as

$$f(x) = 2 \exp\left(-\frac{|x+3|}{2}\right) + \exp(-x^2) + \mathbb{I}_{[2,8]}(x) + \exp(-(x-5)^4),$$

where  $\mathbb{I}_I$  is the indicator function:

$$\mathbb{I}_I(x) = \begin{cases} 1 & \text{if } x \in I \\ 0 & \text{else} \end{cases}.$$

**Question 1** (Density). *Plot the function to get some idea of the targeted distribution.*

**Question 2** (Approximate Monte Carlo estimate). *Under the simplifying assumption that the support of  $f$  is  $[a, b]$  (with well chosen  $a$  and  $b$ ), provide a Monte Carlo estimate of the normalizing constant  $Z_X$ . Study it as a fonction of  $N$ , the number of points to be sampled.*

Listing 1: some useful commands in Python

```
> import numpy as np
> x = np.zeros(10) # Create a numpy array
# of 10 zeros
> x = np.arange(-10, 10, 0.01) # Create a numpy array
# of regularly spaced numbers
> y = 2*np.exp(-np.abs(x+3)/2) + np.exp(-x**2) +
np.where(((x>2)&(x<8)), 1, 0) + np.exp(-abs(x-5)**4)
# numpy processes vectors as Matlab
> import pylab as pl
> pl.plot(x, y)
> pl.show()
> np.random.uniform(a, b, size=4) # Draw 4 samples from Uniform([a, b])
> np.random.uniform() # Draw one sample from Uniform([0, 1])
> np.random.normal(mu, sigma, size=4)
```

```

# Draw 4 samples from Normal(mu, sigma^2)
> np.random.normal() # Draw one sample from Normal(0, 1)
> N, B, _ = pl.hist(samples, bins=k, range=(-15, 15),
                    normed= 1, stacked = 1)
# Draw an histogram of array samples with k bins
# equally distributed on [-15, 15]
# Returns
# * vector N of number of samples in each bin.
# * vector B of X-coordinates of bin separations
> np.mean(x)
# Returns the mean of an numpy array x

```

## Rejection sampling

**Question 3** (Gaussian proposal). *Find a Gaussian density  $q(x)$  and a constant  $M$  such that  $Mq(x) > f(x)$ .*

**Question 4** (Rejection sampling). *Using the rejection sampling algorithm, sample points according to  $f$ :*

1. *check the correctness of the sampled distribution by plotting the histogram;*
2. *how many samples do you waste in average?*
3. *compare this value to the theoretical one (computed using the approximation of  $Z_X$ );*
4. *do the same job with  $M' = 3M$  and comment;*
5. *do the same job with  $M' = \frac{M}{3}$  (assuming the original  $M$  was tight enough, otherwise choose a smaller  $M'$ ) and comment.*

## Markov Chain Monte Carlo

**Question 5** (Metropolis-Hasting). *Using the Metropolis-Hasting algorithm, sample points according to  $f$ , using a Gaussian transition kernel of width  $\sigma \in \{0.02, 1, 50\}$ :*

1. *for each case, show the followed Markov path (sampled points as a function of time);*
2. *check the validity of the sampled distribution by plotting the histogram;*
3. *comment the results.*

## Estimation of a rare event

Now, we would like to estimate the probability of the rare event “ $X \in [15, 20]$ ”, that is  $P(15 < X < 20)$ . In other word, we would like to estimate

$$E[h(X)] \text{ with } h(x) = \mathbb{I}_{[15, 20]}.$$

**Question 6** (Monte Carlo Estimate). *Estimate this probability using a Monte Carlo Estimate (varying the number  $N$  of drawn samples). Comment.*

**Question 7** (Importance sampling with unnormalized distribution). Let  $p$  be the distribution of interest, such that we want to estimate  $\int h(x)p(x)dx$ . Assume that  $\tilde{p}$  is known (up to the normalizing constant). Let  $q$  be the importance proposal. One may also want to use only the unnormalized distribution  $\tilde{q}$ . We recall that the the importance sampling estimator is given by:

$$\frac{1}{N} \sum_{i=1}^N w_i h(x_i) \text{ with } w_i = \frac{p(x_i)}{q(x_i)} \text{ and } x_1, \dots, x_N \sim q.$$

Justify the fact that the following importance sampling estimator is correct :

$$\frac{\frac{1}{N} \sum_{i=1}^N \tilde{w}_i h(x_i)}{\frac{1}{N} \sum_{i=1}^N \tilde{w}_i} \text{ with } \tilde{w}_i = \frac{\tilde{p}(x_i)}{\tilde{q}(x_i)} \text{ and } x_1, \dots, x_N \sim q.$$

Hint: you can use the fact that, writing  $Z_p$  and  $Z_q$  the normalizing constants, we have that  $\frac{Z_p}{Z_q} = \frac{1}{Z_q} \int \tilde{p}(x)dx = \int \frac{\tilde{p}(x)}{\tilde{q}(x)}q(x)dx$ . Be aware of the fact that this requires that  $p(x) > 0 \quad q(x) > 0$ .

**Question 8** (Importance sampling estimate). Use importance sampling to estimate this probability, using as a proposal a Gaussian distribution  $N(\mu, \sigma)$  with  $\mu = 17.5$  and  $\sigma = \{1, 3, 10\}$ :

1. do you obtain the expected result?
2. how do you explain such a bias?

**Question 9** (Importance sampling with a known normalization factor). The aim here is to apply the basic importance sampling estimator:

1. knowing that  $\int_{\mathbb{R}} \exp(-x^4)dx = 2\Gamma(\frac{5}{4}) \approx 1.8128$ , compute the constant  $Z_X$ ;
2. what proposal should you choose here to get a low variance ?
3. estimate the probability of the rare event and compare to previous results.